
Intra-tumoral IL-12 promotes CD8+ T Effector/Memory Cell-mediated, IFN-gamma and FASL dependent elimination of CD4+CD25+Foxp3+ T-suppressor cells from tumors.


Mehmet Okyay Kilinc, Ph.D.
Immunology and Microbiology
School of Medicine, SUNY/Buffalo




Tumors are Infiltrated by CD8+/CD4+ CD25- Tem and CD4+CD25+ T-effector/suppressor Cells




CD8+/CD4+ Tem are Functionally Impaired Whereas CD4+ CD25+ T-cells Demonstrate Suppressive Activity




CD4+ CD25+ Suppression assay







Intra-tumoral IL-12 Induces CD8+ T-cell Activation and Tsuppressor Cell Elimination

Post-Rx CD8+ T-cell Activation






Post-Rx T-cell kinetics




Post-Rx Foxp3+ Tsuppressor kinetics




Tsupp Loss is Associated with Apoptotic Cell Death

Annexin V binding to Tsupp



IL-12-induced T-suppressor Cell Apoptosis is CD8+ T-cell, IFN_γ and FasL-dependent



Summary

- Tumors are infiltrated by functionally-impaired CD8/CD4 Tem and fully-active Tsupp.
- Intra-tumoral IL-12 induces the activation of preexisting Tem, elimination of Tsupp and infiltration of CD8+ Teff.
- Elimination of Tsupp is due to apoptotic death and is dependent on pre-existing CD8+ Tem, IFN_γ and FasL.



I would like to thank to P.I of the lab; Nejat Egilmez (Associate Prof.) and other lab members; Tao Gu (Post- doc), Rachael Rowswell (Ph.D student), Jamie Harden (Ph.D student), Lauren Virtuoso (Lab tech.), Maryann Hochberg (Lab tech.) and Wyeth Pharmaceuticals.

- -Mehmet Okyay Kilinc (Res. Asst. Prof.)
- -E-mail:mokilinc@buffalo.edu