

## Presenter disclosures information David H. Munn, MD

In compliance with ACCME policy, the following are disclosed to the activity audience:

Research Support	NewLink Genetics, Inc.
Employee	Medical College of GA (intellectual property)
Consultant	NewLink Genetics, Inc.
Scientific Advisory Board	NewLink Genetics, Inc.

# Immunotherapy of Cancer: Statement of the problem

 Immunotherapy of cancer must do more than simply present antigens to the immune system

... it must disrupt a pre-existing state of functional tolerance toward tumor antigens

### Tumor-induced tolerance

### Tolerance is <u>acquired</u>, <u>active</u> and <u>dominant</u>

- <u>acquired</u> because even truly foreign antigens will become tolerated if introduced on tumors
- <u>active</u> because tolerance cannot be overcome simply by a good antigen and a strong adjuvant
  - .... (i.e., responses are actively suppressed)
- <u>dominant</u> because even vaccination or adoptive transfer of pre-activated T cells is still subject to suppression

### Indoleamine 2,3-dioxygenase (IDO)

- IDO is a natural endogenous molecular mechanism of immune suppression
  - involved in pregnancy, mucosal tolerance
- IDO can create acquired peripheral tolerance de novo

## Haplo-mismatched allografts transfected with IDO are tolerated without additional immunosuppression





figure adapted from KA Swanson, David S. Wilkes et al

Am. J. Resp. Cell Molec. Biol. Vol. 30, pp. 311-318, 2004 © 2004 American Thoracic Society


### IDO and malignancy


- IDO is expressed by cancer cells in a range of tumor types
- High IDO expression appears correlate with poor outcome in a number of cancers
  - ovarian cancer
  - AML
  - endometrial carcinoma
  - colon cancer
  - melanoma



## Predictive value of abnormal IDO expression in human tumor-draining lymph nodes

- 40 patients with cutaneous malignant melanoma, no metastases
- sentinel lymph node obtained at time of initial diagnosis
- in collaboration with Scott Antonia at Moffitt Cancer Center





#### IDO-inhibitor (NSC-721782):

1-methyl-[D]-tryptophan (D-1MT)

### 1MT is synergistic with chemotherapy

(MMTV-neu tumor model – Prendergast lab)

Table 1 IDO inhibition enhances the efficacy of certain commonly used cancer chemotherapeutic agents


Compound	Class	Mean ± s.e. (+1MT)	Mean ± s.e. (–1MT)	P	n	Dose (mg/kg)	Route	Schedule
Cisplatin	Alkylating agent	0.77 ± 0.18	1.7 ± 0.33	0.0419	7,8	1.0	i.v.	3×/week
Cyclophosphamide	Alkylating agent	$0.81\pm0.12$	$1.4\pm0.18$	0.0269	5,5	100	i.v.	3x/week
Doxorubicin	Antineoplastic antibiotic	0.79 ± 0.07	$1.5\pm0.25$	0.0150	6,4	0.66	i.v.	3x/week
5-Fluorouracil	Antimetabolite	1.2 ± 0.20	1.1 ± 0.25	0.8926	8,7	50	i.v.	3×/week
Methotrexate	Antimetabolite	$1.7 \pm 0.28$	$1.7\pm0.38$	0.9047	3,3	1.0	i.v.	3×/week
Paclitaxel	Mitotic inhibitor (taxane)	0.68 ± 0.11	2.4 ± 0.43	0.0010	8,7	13.3	1.v.	3x/week
Vinblastine	Mitotic inhibitor (vinca alkaloid)	$1.3\pm0.19$	$1.2\pm0.18$	0.7368	10,8	1.0	i.v.	3x/week
FTI	Signal transduction inhibitor	0.67 ± 0.11	1.0 ± 0.16	0.0979	8,8	40	i.p.	qdx11
Rapamycin	Signal transduction inhibitor	$0.97 \pm 0.07$	$0.99 \pm 0.25$	0.9417	4,4	1.5	i.v.	qdx11
Tetrathiomolybdate	Antiangiogenic (iron chelator)	$1.9 \pm 0.52$	$2.0\pm0.42$	0.7996	3,4	40	p.o.	qdx11
Vehicle		1.7 ± 0.17	3.0 ± 0.44	0.0061	12,5			

from Muller et al

NATURE MEDICINE VOLUME 11 NUMBER 3 MARCH 2005


Take-home message:

even genetically-diverse, spontaneous tumors show widespread dependence on the IDO mechanism after chemoRx




D vs L isomer of 1MT (4T1 breast-tumor model) (courtesy of George Prendergast lab, from Hou et al, Cancer Res. 2007)

### IDO and Tregs form a mutually-reinforcing system




### IDO directly activates (Tregs) in TDLNs


(Sharma et al, J. Clin. Invest., 2007)




From Sharma et al Blood 113:6102, 2009




IDO-inhibitor (1MT) is synergistic with vaccine against established tumors




# IDO regulates re-programming of Tregs into TH17-like T-helper cells in tumor-draining lymph nodes





## B DCs in TDLN after vaccination



IDO regulates Treg conversion to TH17 cells in part via CGN2-mediated suppression of IL-6 expression

www.asco.org

# Phase I Trial of 1-methyl-D-tryptophan

PI: Scott Antonia MD PhD

Co PI: Hatem Soliman MD

Dan Sullivan MD

MOFFITT CANCER CENTER

Moffitt Cancer Center/Southeast Phase II Consortium

Chuck Link MD Nick Vanahanian MD William Ramsey MD PhD

NewLink Genetics Inc

# Combination of IDO-inhibitor drugs with chemotherapy and immunotherapy

#### Chemotherapy

- IDO appears to be a non-redundant mechanism needed by the tumor to <u>re-establish</u> the suppressive milieu following chemotherapy
- blocking IDO may thus promote immune response against the tumor after chemotherapy

### **Immunotherapy**

- Tumor-induced IDO acts as a fundamental antagonist to anti-tumor immune responses generated by immunotherapy
- Blocking IDO allows re-programming of Tregs into TH17-like helper cells following vaccination

### Lessons and Take Home Messages

#### Key points

- IDO acts to create a suppressive milieu in tumor and tumordraining LN
- IDO <u>promotes</u> Treg activation, and <u>prevents</u> vaccine-induced reprogramming of Tregs into T-helper cells

### Potential impact on the field

- IDO-inhibitor drugs can be synergistic with chemotherapy, and together may open a "window of opportunity" for vaccines and other immunotherapy
- The combination of a vaccine plus an IDO-inhibitor drug may be able to re-program Tregs in situ into non-suppressive T-helper cells (which may offer an alternative to Treg depletion)

### Acknowledgements

- Andrew Mellor
- Madhav Sharma
- Deyan Hou

Medical College of Georgia

- Scott Antonia
- Hatem Soliman

Moffitt Cancer Center

- George Prendergast
- Alex Muller

Lankenau Institute