

Immunotherapy for the Treatment of Lung Cancer

Clarke Low, MD

Director of Thoracic Oncology
Intermountain Healthcare


Disclosures

- No disclosures to report
- I will not be discussing non-FDA approved indications during my presentation.

Immunotherapy for the Treatment of Lung Cancer

Checkpoint Inhibitors: PD-1 and PD-L1


- PD-1 acts as an “off-switch” for T cells when interacting with PD-L1
- Tumor PD-L1 expression allowing cancer cells to evade immune attack
- Antibodies against PD-1 and PD-L1 boost the immune response against cancer cells



Gong J, Journal for Immunotherapy of Cancer, 2018


Combination Immune Checkpoint Blockade

- CTLA-4 acts as an “off-switch” for T cells when interacting with B7
- Combination strategies combine both CTLA-4 and PD-1/PD-L1 blockade



Ribas A, NEJM, 2012

FDA-approved Checkpoint Inhibitors in NSCLC




CA209-003: Nivolumab in Heavily-pretreated Advanced NSCLC (NCT00730639)

Phase 1, 5-Year Update

5-Year Survival

- First report of long-term survival rate in patients with metastatic NSCLC treated with an immune checkpoint inhibitor
- According to the National Cancer Institute's SEER data, 5-year survival rate for patients with advanced NSCLC is 4.9%



Gettinger et al. JCO 2018
Brahmer et al, AACR 2017
NCI SEER data, Lung and Bronchus Cancer, 2014


Treatment Naïve Regimens: Competing Strategies

- KEYNOTE 024 – Pembrolizumab vs. Chemotherapy in PD-L1 > 50%
- KEYNOTE 042 – Pembrolizumab vs. Chemotherapy in PD-L1 > 1%
- KEYNOTE 189 – Pembrolizumab + Chemotherapy vs. Chemotherapy alone in patients with advanced non-squamous NSCLC
- IMPOWER 150 – Atezolizumab + Chemotherapy (Bev) vs. Chemotherapy (Bev) in patients in advanced non-squamous NSCLC
- KEYNOTE 407 – Pembrolizumab + Chemotherapy vs. Chemotherapy in advanced squamous cell lung cancer
- Checkmate 227 – Ipilimumab + Nivolumab vs. Chemotherapy in advanced NSCLC with high TMB

KEYNOTE-024: Pembrolizumab vs. Chemotherapy for PD-L1 Positive (>50%) NSCLS Study Design (NCT021427389)


Key Eligibility Criteria

- **Untreated** stage IV NSCLC
- PD-L1 TPS $\geq 50\%$
- ECOG PS 0-1
- No activating *EGFR* mutation or *ALK* translocation
- No untreated brain metastases
- No active autoimmune disease requiring systemic therapy



Reck M et al, ESMO 2016, NEJM 2016

KEYNOTE-024: Pembrolizumab vs. Chemotherapy for PD-L1 >50% NSCLC Overall Survival



Reck M et al, ESMO 2016, NEJM 2016

KEYNOTE-042: Pembrolizumab vs. Chemotherapy for PD-L1 > 1% NSCLC

KEYNOTE-042 Study Design

Key Eligibility Criteria

- Untreated locally advanced or metastatic NSCLC of any histology
- PD-L1 TPS $\geq 1\%$
- No sensitizing *EGFR* or *ALK* alterations
- ECOG PS 0 or 1
- No untreated or unstable CNS metastases
- No history of pneumonitis that required systemic corticosteroids

Stratification Factors

- Region (east Asia vs rest of the world)
- ECOG PS (0 vs 1)
- Histology (squamous vs nonsquamous)
- PD-L1 TPS ($\geq 50\%$ vs 1-49%)

Randomize
1:1

N = 637

Pembrolizumab
200 mg Q3W
for up to 35 cycles

N = 637

Carboplatin AUC 5 or 6 Q3W +
Paclitaxel 200 mg/m² Q3W^a
OR
Carboplatin AUC 5 or 6 Q3W +
Pemetrexed 500 mg/m² Q3W^a
for up to 6 cycles

End points


- Primary: OS in PD-L1 TPS $\geq 50\%$, $\geq 20\%$, and $\geq 1\%$
- Secondary: PFS and ORR in TPS $\geq 50\%$, $\geq 20\%$, and $\geq 1\%$; safety in TPS $\geq 1\%$

^aPemetrexed maintenance therapy was optional but strongly encouraged for patients with nonsquamous histology.


Lopes et al, ASCO 2018

KEYNOTE-042: Pembrolizumab vs. Chemotherapy for PD-L1 > 1% NSCLC Overall Survival

Overall Survival: TPS ≥ 50%




Overall Survival: TPS ≥ 1-49% (Exploratory Analysis^a)



Survival benefit seemed to be driven by the TPS > 50% subset with little benefit witnessed in the subset TPS > 1- 49%


Lopes et al, ASCO 2018

KEYNOTE-189: Pembrolizumab/Carboplatin /Pemetrexed vs Chemotherapy for Advanced Non-squamous NSCLC



Ghandi et al, NEJM 2018


KEYNOTE-189: Pembrolizumab/Carboplatin /Pemetrexed vs Chemotherapy for Advanced Non-squamous NSCLC




Ghandi et al, NEJM 2018

KEYNOTE-189: Pembrolizumab/Carboplatin /Pemetrexed vs Chemotherapy for Advanced Non-squamous NSCLC


PD-L1 < 1%



PD-L1 1 – 49%




PD-L1 ≥ 50%



Ghandi et al, NEJM 2018


KEYNOTE-407: Pembrolizumab/chemotherapy vs Chemotherapy for Advanced Squamous-cell NSCLC




Paz-Ares et al, ASCO 2018

KEYNOTE-407: Pembrolizumab/chemotherapy vs Chemotherapy for Advanced Squamous-cell NSCLC

PFS (RECISTv1.1, BICR)




Overall Survival



Paz-Ares et al, ASCO 2018

IMPOWER 150: Atezolizumab/Carboplatin/ Paclitaxel/Bevacizumab vs Carboplatin/ Paclitaxel/ Bevacizumab in advanced non-squamous NSCLC




Socinski et al, NEJM 2018


IMPOWER 150: Atezolizumab/Carboplatin/ Paclitaxel/Bevacizumab vs Carboplatin/ Paclitaxel/Bevacizumab in advanced non-squamous NSCLC

Landmark OS, %	Arm B: atezo + bev + CP	Arm C: bev + CP
12-month	67%	61%
18-month	53%	41%
24-month	43%	34%

HR^a, 0.78
 (95% CI: 0.64, 0.96)
P = 0.0164
 Median follow-up: ~20 mo




Subgroup	n (%) ^a
PD-L1–High (TC3 or IC3) WT	136 (20%)
PD-L1–Low (TC1/2 or IC1/2) WT	226 (32%)
PD-L1–Negative (TC0 and IC0) WT	339 (49%)
Liver Metastases WT	94 (14%)
No Liver Metastases WT	602 (86%)
ITT (including EGFR/ALK+)	800 (100%)
EGFR/ALK+ only	104 ^b (13%)
ITT-WT	696 (87%)




Socinski et al, NEJM 2018

Tumor Mutational Burden (TMB) may Determine Sensitivity to PD-1 Blockade in NSCLC

- In two independent cohorts, higher nonsynonymous tumor mutational burden (TMB) was associated with improved objective response, durable clinical benefit, and PFS.




*Partial or stable response lasting > 6 mo



Rizvi N et al, Science, 2015

CheckMate 227: Ipilimumab + Nivolumab vs Chemotherapy in TMB-high patients





Hellman et al, NEJM, 2018

CheckMate 227: Ipilimumab + Nivolumab vs Chemotherapy in TMB-high patients

Hazard ratio for disease progression or death,
0.58 (97.5% CI, 0.41–0.81)
P<0.001

Patients with Progression-free
Survival (%)





Median Duration
(95% CI)
mo

NR (12.2–NR)
5.4 (4.2–6.9)

Hellman et al, NEJM, 2018

CheckMate 227: Ipilimumab + Nivolumab vs Chemotherapy in TMB-high patients



Hellman et al, NEJM, 2018

PD1/PD-L1 Inhibitors Increase *Overall Survival* in 2L Advanced NSCLC

CHECKMATE 017 (nivolumab)

	Median Overall Survival mo (95% CI)	1-Yr Overall Survival % of patients (95% CI)	No. of Deaths
Nivolumab (N=135)	9.2 (7.3–13.3)	42 (34–50)	86
Docetaxel (N=137)	6.0 (5.1–7.3)	24 (17–31)	113

CHECKMATE 057 (nivolumab)

	Nivolumab (n = 292)	Docetaxel (n = 290)
mOS, mo	12.2	9.4
HR = 0.73 (96% CI: 0.59, 0.89); P = 0.0015		


KEYNOTE 010 (TPS ≥ 1%) (pembrolizumab)

Treatment Arm	Median (95% CI), mo	HR* (95% CI)	P
Pembro 2 mg/kg	14.9 (10.4-NR)	0.54 (0.38-0.77)	0.0002
Pembro 10 mg/kg	17.3 (11.8-NR)	0.50 (0.36-0.70)	<0.0001
Docetaxel	8.2 (6.4-10.7)	--	--

OAK (atezolizumab)


HR, 0.73^a
 (95% CI, 0.62, 0.87)
 P = 0.0003
Minimum follow up = 19 months

PACIFIC (NCT02125461): Durvalumab after Chemoradiotherapy in Stage III NSCLC



1. In House Data, AstraZeneca Pharmaceuticals LP. PACIFIC Protocol. 2014.
2. NIH 2015 NCT02125461, <http://clinicaltrials.gov/ct2/show/NCT02125461>.
3. Creelan B, Iannotti NO, Salamat MA, et al. 2016. (PHRR150325-000989)
4. Ann Oncol. 2015;26 (supplement 1): i24-i28, abstract 95TIP.



PACIFIC (NCT02125461): Durvalumab after Chemoradiotherapy in Stage III NSCLC



Antonia et al, NEJM 2017

Checkpoint Inhibitors in Metastatic EGFR-Mutated NSCLC

Meta-Analysis: CM-057, KN-010, POPLAR




CK Lee et al., JTO 2016

Single-agent Toxicities in 2/3L Randomized Trials

	Atezolizumab OAK	Nivolumab SQ: CM 017 (updated OS; 2L)	Nivolumab NSQ:CM 057 (updated OS; 2/3L)	Pembrolizumab Keynote 010
Related Grade 3-5 AEs	15%	8%	11%	13-16%
Discontinuation due to related AEs	5%	6%	6%	4-5%
Pneumonitis AEs	1%	5%	3%	4-5%


Rittmeyer, et al., *Lancet* 2017
 Brahmer, et al., *NEJM* 2015
 Borghaei, et al., *NEJM* 2015
 Herbst, et al., *Lancet* 2015

KEYNOTE-189: Pembrolizumab/Carboplatin /Pemetrexed vs Chemotherapy for Advanced Non-squamous NSCLC



Ghandi et al, NEJM 2018

KEYNOTE-407: Pembrolizumab/chemotherapy vs Chemotherapy for Advanced Squamous-cell NSCLC



Paz-Arez et al, ASCO, 2018

CheckMate 227: Ipilimumab + Nivolumab vs Chemotherapy in TMB-high patients

TRAE, ^a %	Nivolumab + ipilimumab (n = 576)		Chemotherapy (n = 570)	
	Any grade	Grade 3–4	Any grade	Grade 3–4
Any TRAE	75	31	81	36
TRAE leading to discontinuation^b	17	12	9	5
Most frequent TRAEs (≥15%)				
Rash	17	2	5	0
Diarrhea	16	2	10	1
Fatigue	13	1	18	1
Decreased appetite	13	<1	19	1
Nausea	10	<1	36	2
Constipation	4	0	15	<1
Anemia	4	2	32	11
Neutropenia	<1	0	17	9
Treatment-related deaths^c	1		1	

Hellman et al, NEJM, 2018

Summary of Frontline Strategies in Advanced NSCLC



Clinical Trial	Drug	PFS (Months)	OS (Months)	PFS HR in PD-L1 neg	Toxicities Grade 3 - 5
KEYNOTE-024 PD-L1 ≥ 50%	Pembro	10.3	30	NA	31% vs 53%
	Plat/Pem or Gem or Pacli	6	14.2		
KEYNOTE-042 PD-L1 ≥ 1%	Pembro	5.4	16.7	NA	18% vs 41%
	Plat/Pem or Pacli	6.5	12.1		
IMpower150 Non-squamous	Atezo + Beva + Carbo/Pacli	8.3	19.2	0.77	60 vs 51%
	Beva + Carbo/Pacli	6.8	14.7		
KEYNOTE-189 Non-squamous	Pembro + Plat/Pem	8.8	NR	0.75	67% vs 66%
	Plat/Pem	4.9	11.3		
KEYNOTE-407 Squamous	Pembro + Carbo/Pacli or NabPacli	6.4	15.9	0.68	70% vs 68%
	Carbo/Pacli or NabPacli	4.8	11.3		
CheckMate 227 TMB≥10mut/Mb	Nivo + Ipi	7.2	23	0.48	31% vs 36%
	Plat/Pem or Gem	5.4	16.7		

Adapted from Solange Peters, 2018 ASCO Annual Meeting * This is for illustration purposes only and comparing different trials is challenging as populations, indications, and other characteristics vary.


Case Study 1

- 69 y/o man, retired woodworker with 50 pack year history
- History of latent TB that was treated, BPH, remote methamphetamine abuse
- Presents with 2 month history of dysphonia, productive cough
- CXR demonstrates ~4cm LLL mass
- CT Chest: 4.8x3.2x7.2 cm lingular mass; prominent L hilar LN
- Staging PET/CT: liver and boney metastases
- Staging MRI Brain: calvarial and C1/C2 mets
- Percutaneous core lung biopsy: adenocarcinoma, TTF-1+
 - EGFR, ALK, ROS1 negative
 - PD-L1 81-90%

At diagnosis



6 months on treatment



Case Study 1

- 7 months on treatment; Tbili 1.4, AST/ALT 149/232
 - Grade 2 Hepatitis
 - How would you manage?
 - Stopped pembrolizumab; started prednisone 1mg/kg
- 1 month later: LFT's have all normalized; start to taper
- 2 more months: patient self-discontinued steroids; Tbili 10.7, AST/ALT 105/144 (Grade 3 hepatitis; 3-10x ULN Tbili)
- 5 more months of steroids with slow taper

Case Study 1

- 11 months after stopping pembrolizumab has asymptomatic radiographic progression
- Alive 24 months after Stage IV NSCLC diagnosis w/ good QoL on maintenance pemetrexed

Case study 2

- 76 y/o man, retired software engineer, former smoker w/ 40 pack year hx
- History of HTN, HLD, prediabetes
- Presents with 2 month history of progressive DOE
- CXR demonstrates moderate L pleural effusion
- Thoracentesis: lung adenocarcinoma, TTF-1 +; insufficient material for more testing
- Endobronchial biopsy: poorly differentiated adenocarcinoma
 - EGFR, ALK, ROS1, BRAF negative
 - PD-L1 1%
- No distant sites of metastasis; Stage IVA

Case study 2

- How would you treat?
 - Carboplatin and pemetrexed
 - Carboplatin and pemetrexed and pembrolizumab
 - Pembrolizumab monotherapy

Case study 2

- Started carboplatin / pemetrexed / pembrolizumab
 - After one cycle develops grade 1 dermatitis of forearms
 - After two cycles has stable disease radiographically
 - After three cycles develops grade 3 pancytopenia, ANC 1400, Hb 5.2, Plt 20
 - Has been receiving IM B12 and taking folic acid
 - No evidence of hemolysis, B12 and folate levels normal
 - Supportive transfusions and cytopenias resolve within 1-2 weeks
- Discontinue chemotherapy and start maintenance pembrolizumab

Case study 2

- 5 months after starting treatment
 - Clinical progression (worsening dyspnea on exertion, chest pain, fatigue)
 - Radiographically stable disease, no findings of pneumonitis
 - Patient elects to go onto hospice; dies ~4 weeks later